Patient Testimonial

MARIUS PHARMACEUTICALS

CONSENT, RELEASE OF CLAIMS, AND WAIVER OF LIABILITY

I wish to provide my experiences as a patient as part of MARIUS's product and/or disease awareness campaigns ("Compaigns"). I understand that my participation in the Campaigns is completely voluntary, without compensation, and at my own risk.

By signing this Consent and Release ("Agreement"), I hereby grant to MARIUS and its affiliates, successors, and assigns (collectively, "MARIUS") the limited permission to photograph me, copyright, use, and publish printed, video, audio or photographic images of me and/or my statements, including images in which I may be included, in whole or in part, regarding my experience as a patient with low testosterone who is prescribed KYZATREX™ (testosterone undecanoate) for treatment (collectively, "Content"). MARIUS may reproduce, publish, display, post, or otherwise use the Content in all media, including but not limited to social media, marketing materials, websites, and online, for a period not to exceed five (5) years from the Effective Date of this Agreement. In all cases, Content shall be used solely for the purpose of publicity, advertising, marketing, and promotion in relation to the Campaigns.

I understand that MARIUS may use the Content by itself or combine it with other written, printed, graphic, or audio matter. MARIUS has the right to edit, abridge, and format the Content, in whole or in part, in all media now known or later developed for the Campaigns, and I waive all rights to inspect or approve the finished products. MARIUS may contact me to ensure that Content accurately reflects my personal experience with low testosterone and/or KYZATREX.

I hereby grant MARIUS full ownership of any such Content subject to the use restrictions of this Agreement. I waive all rights that I may have to claims for payment or royalties in connection with any exhibition, posting, or publication of the Content. I hereby waive any and all claims that I may have against MARIUS, and will not hold MARIUS responsible for any claim or liability, related to their use of the Content so long as that use is consistent with the terms of this Agreement, including but not limited to claims in the nature of copyright infringement, trademark infringement, defamation, disparagement, libel, slander, invasion of privacy or publicity or the like. I understand and acknowledge that MARIUS: (a) will be under no obligation to actually use the Content, and (b) has the right to use the Content either with or without my name. Any use of my name will be limited to my first name or my first name and last name initial.

By signing below, I warrant that all information I submit to MARIUS, including the Content, is true and accurate to the best of my knowledge. I have the right to grant all necessary permissions for MARIUS's use of the Content; I am at least 18 years of age, over the age of majority in my jurisdiction, and able to enter into this Agreement. I further acknowledge that I have read and agree to the terms of this Agreement. This Agreement shall be binding up on me and my heirs, legal representatives, and assigns. This Agreement is governed by the laws of North Carolina, and any actions that in any way arise from or are related to the Content or Campaigns must be brought in the state or federal courts located in North Carolina.

Download PDF

Important Safety Information for KYZATREX™ (testosterone undecanoate)

KYZATREX™ can increase blood pressure, which can increase the risk of having a heart attack or stroke and can increase risk of death due to a heart attack or stroke. Your risk may be greater if you have already had a heart attack or stroke or if you have other risk factors for heart attack or stroke.

Printed Full Name *

Email Address *

Phone *

Street Address *

Apartment, suite, etc.

City *

State/Province *

ZIP / Postal Code *

Date Signed *
01/22/2024

Signature *

Patient Testimonial

First Name *

Last Initial *

Age *

How long have you been taking KYZATREX® (testosterone undecanoate) capsules? *

What is your current dose of KYZATREX? *

Were you previously on testosterone therapy or are you new to testosterone therapy? *
Please Select.

If you are new to testosterone therapy, please select the symptoms that led you to consult your doctor? (Please select all that apply.) *

- Low libido (sex drive)
- Reduced erectile function
- Fatigue/low energy
- Decreased muscle mass
- Weight gain
- Hair loss
- Mood changes
- Reduced bone density
- Anemia (low red blood cell count)
- Other
- I did not have any symptoms
- I am not new to testosterone therapy

If you were previously on testosterone therapy, why did you switch to a new therapy? (Please select all that apply.) *

- Financial reasons
- Accessibility reasons
- I experienced unwanted side effects.
- I didn't like the administration method or frequency of my old therapy.
- My doctor recommended the switch.
- I am new to testosterone therapy.
- Other

What made you choose KYZATREX over other testosterone replacement therapies? *

- Cost
- Accessibility
- Route of administration
- Doctor recommendation
- Recommendations from family/friends
- Tolerability/side effect profile
- Formulation
- Other

Based on your most recent lab results, what are your current total serum testosterone levels?

How has the introduction of KYZATREX into your routine affected your daily activities and lifestyle?

In what ways has KYZATREX impacted your overall quality of life?

Is there anything else you'd like to share about your experience with KYZATREX?

Submit

Copyright © 2023 KYZATREX™, all rights reserved.
This site is intended for U.S. residents only.

Use

KYZATREX™ (testosterone undecanoate) is a prescription drug that is used to treat adult men who have low or no testosterone levels due to certain medical conditions.

KYZATREX™ is a controlled substance (CIII) because it contains testosterone. It is not known if KYZATREX™ is safe or effective in males younger than 18 years old. Improper use may affect bone growth in children. KYZATREX™ is not meant for use by women.

Important Safety Information for KYZATREX™

KYZATREX™ can increase blood pressure, which can increase the risk of having a heart attack or stroke and can increase risk of death due to a heart attack or stroke. Your risk may be greater if you have already had a heart attack or stroke or if you have other risk factors for heart attack or stroke.

1. If your blood pressure increases while on KYZATREX™, blood pressure medicines may need to be started. If you are currently taking blood pressure medicines, they may need to be changed or new blood pressure medicines may need to be added to control your blood pressure.
2. If your blood pressure cannot be controlled, KYZATREX™ may need to be stopped.
3. Your healthcare provider will monitor your blood pressure while you are being treated with KYZATREX™.

Do not take KYZATREX™ if you: have breast cancer; have or might have prostate cancer; are a woman who is pregnant (KYZATREX™ may harm your unborn baby); are allergic to KYZATREX™ or any of its ingredients; or have low testosterone without certain medical conditions (e.g., do not take KYZATREX™ if you have low testosterone due to age).

Before you take KYZATREX™, tell your healthcare provider about all of your medical conditions, including if you: have high blood pressure or are treated for high blood pressure; have a history of diabetes; have heart problems; have high red blood cell count (hematocrit) or high hemoglobin laboratory value; have urinary problems due to an enlarged prostate; have liver or kidney problems; or have problems breathing while you sleep (sleep apnea).

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking KYZATREX™ with certain other medicines can affect each other. **Especially, tell your healthcare provider if you take:** insulin; medicines that decrease blood clotting (blood thinners); corticosteroids; or medicines that increase blood pressure, such as some cold medicine and pain medicines.

KYZATREX™ may cause other serious side effects including:

1. **Increase in red blood cell count (hematocrit) or hemoglobin,** which can increase the risk of blood clots, strokes, and heart attacks. You may need to stop KYZATREX™ if your red blood cell count increases.
2. **If you already have an enlarged prostate, your signs and symptoms may worsen while taking KYZATREX™.** These may include: increased urination at night; trouble starting your urine stream; urinating many times during the day; urge to go to the bathroom right away; a urine accident; inability to pass urine or weak urine flow.
3. **Increased risk of prostate cancer.**
4. **Blood clots in the legs or lungs.** Signs and symptoms of a blood clot in your leg can include pain, swelling or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain.
5. **Abuse.** Testosterone can be abused when taken at higher than prescribed doses and when used with other anabolic androgenic steroids. Abuse can cause serious heart and psychological side effects.
6. **In large doses, KYZATREX™ may lower your sperm count.**
7. **Liver problems.** Symptoms of liver problems may include: nausea or vomiting; yellowing of your skin or whites of your eyes; dark urine; pain on the right side of your stomach area (abdominal pain).
8. **Swelling of your ankles, feet, or body (edema), with or without heart failure.**
9. **Enlarged or painful breasts.**
10. **Breathing problems while you sleep (sleep apnea).**

Call your healthcare provider right away if you have any of the serious side effects listed above.

The most common side effect of KYZATREX™ is high blood pressure. Other side effects may include: headache, joint or back pain, diarrhea, increased red blood cell count, anxiety, constipation, swelling of the legs, and increased prostate specific antigen (PSA) levels.

These are not all the possible side effects of KYZATREX™. For more information, ask your healthcare provider or pharmacist.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to Marius by visiting www.mariuspharma.com.

Keep KYZATREX™ and all medicines out of the reach of children.

See Full Prescribing Information and Medication Guide for KYZATREX™.